

Table 1

***Wolbachia* prevalence rates in local populations of corn borer of the *Ostrinia* genus**

Insect sampling site	Forage plant	Examined insect number		Prevalence rate, % \pm SE
		total	infected	
Slavyansk-na-Kubani town, Slavyansk District, Krasnodar Territory	<i>Zea</i> ¹	30	2	6.7 \pm 4.6
	<i>Artemisia</i> ²	25	3	12.0 \pm 6.5
Slobodka settlement, Slavyansk District, Krasnodar Territory	<i>Zea</i> ¹	34	1	2.9 \pm 2.9
	<i>Artemisia</i> ²	21	3	14.3 \pm 7.7
	<i>Cannabis</i> ²	38	25	65.8 \pm 7.7
Botanika settlement, Gulkevichi District, Krasnodar Territory	<i>Zea</i> ¹	34	7	20.6 \pm 7.0
	<i>Cannabis</i> ²	38	6	15.8 \pm 6.0
Maiskiy settlement, Belgorod District, Belgorod Region	<i>Zea</i> ¹	43	9	20.9 \pm 6.3
	<i>Artemisia</i> ²	16	7	43.8 \pm 12.5

¹*Ostrinia nubilalis*; ²*Ostrinia scapularis*; SE – standard error