

Mamayeva A.V., Yaschuk A.G.

Bashkir State Medical University, Russia

THE SEARCH FOR THE TECHNIQUE OF HYSTERECTOMY IN FEMALES WITH CONNECTIVE TISSUE DYSPLASIA

Purpose. To determine differential etiopathogenic approach of hysterectomy in females to prevent the occurrence of genital prolapse after hysterectomy.

Materials and methods. It has been proved that systemic connective tissue defect may result in the development vaginal and uterine prolapse, elyptosis and descent of the womb alongside with the occurrence of such pathologies as hernias, splanchnoptosis, varicose disease, hemorrhagic diatheses. We performed complex examination of 210 females.

Results. We identified a group of 37 patients who underwent hysterectomy and had manifestations of connective tissue dysplasia (CTD) in the form of genital prolapse. The onset of genital ptosis (descent) and its aggravation were already observed during the reproductive period as the major pathology in 15 females, after uterine amputation – in 18 females and after transabdominal uterine extirpation – in 4 females. These patients underwent transvaginal uterine extirpation (13 cases), transvaginal extirpation of uterine cervix stump (18), and anterior and posterior colporrhaphy combined with perineolevatoroplastics (6 cases). Recurrences of genital prolapse were revealed postoperatively in 8 cases: within the first year – in 1 female, within 2-4 years – in 6 females, after 15 years – in 1 subject. Genital prolapses occurring after transvaginal hysterectomy were found in 3 cases, after anterior colporrhaphy and colpoperineolevatoroplasty in 4 females, after transvaginal extirpation of the cervical stump, anterior colporrhaphy

and colpoperineolevatoroplasty in 1 female. Different manifestations of CTD (in combination from 3 to 10 symptoms and more in each patient) were revealed in all patients. When assessing values of hemostasiogram all patients were found to demonstrate thrombocytopenia, in 26 patients the decrease of collagen-induced and ADF-induced thrombocyte aggregation were revealed. The activity of von Willebrand's factor was markedly reduced in 18 patients, hyperfibrinogenemia was found in 11 patients. Oxyproline excretion in the daily urine test increased two-fold and more compared to normal levels in 29 females. Serum magnesium levels corresponded either to the lowest values or were below normal values. According to the grading scheme of phenotypic and clinical picture of CTD severe forms were revealed in 5 patients, moderately severe forms in 19 patients and mild forms in 13 patients.

Conclusions. The most frequent cause of genital prolapse as well as its recurrence that occur after transvaginal hysterectomy, abdominal extirpation and uterine amputation is the connective tissue defect. To prevent genital prolapse in patients with CTD it is advisable to perform transvaginal hysterectomy with sacro-spinal MESH-vaginopexy. In abdominal hysterectomy it is recommended to perform sacrovaginopexy in such patients to avoid genital prolapse. In recurrent cases of vaginal stump prolapse (uterine cervix) sacro-spinal MESH-vaginoplasty with simultaneous transvaginal stump extirpation (uterine cervix) is indicated.

Plekhanov A.N.

City center of laparoscopic surgery of the Hospital by St.Elizabeth, St.Petersburg, Russia

ADVANTAGES OF LAPAROSCOPIC-ASSISTED VAGINAL HYSTERECTOMY OVER ABDOMINAL HYSTERECTOMY IN PATIENTS WITH FIBROIDS

Subject matter. Subject matter was the comparative study of laparoscopic-assisted vaginal hysterectomy (LAVH) and abdominal hysterectomy (AH) in patients with 12-22 weeks pregnancy fibroid.

Material and methods. The results of comparative study of laparoscopic-assisted vaginal hysterectomy (134 cases) and abdominal hysterectomy (109 cases) in the patients with a big size fibroid in 2000-2004 yy are presented.

Results. Time of operation was $67 \pm 6,3$ min (AH),

$87,21 \pm 5,6$ min (LAVH), blood loss was $315 \pm 9,6$ ml (AH), $250 \pm 17,1$ ml (LAVH), weight of fibroid was $712 \pm 8,2$ g (AH) and $704,5 \pm 9,25$ g (LAVH).

Conclusion. Laparoscopic-assisted vaginal hysterectomy has the following advantages: minimal traumatization, favorable cosmetic effect, short postoperative and rehabilitation period, absence of complications in the late postoperative period, decrease of the risk of intraoperative complications. The experience accumulated proves the LAVH technique to be reasonable and efficient.