

Povzun S.A.¹, Bezhenar V.F.^{2,3}, Tsvelev Iu.V.²,
Fridman D.B.^{1,2}

¹ Department of Pathomorphology,

² Department of Obstetrics and Gynecology Military- medical academy,

³ The D.O.Ott Research Institution of Obstetrics and Gynecology RAMS, Saint Petersburg, Russia.

EFFICIENCY OF DIFFERENT BIOPSY METHODS OF ADENOMYOSIS DIAGNOSIS, COMPARATIVE ANALYSIS

Introduction. Usage of low invasive methods in adenomyosis diagnosis helps to diagnose adenomyosis definitely by histological examination of myometrium samples.

Material and methods. Hysterectomy specimen (n=32), in 24 (75%) cases the adenomyosis was confirmed by pathologic examination. Imitation of transcervical and transabdominal puncture biopsy, pinch and resection biopsy in vitro were performed.

Results:

- pinch biopsy is characterized by low volume of myometrium sample ~1mm³, unavailability to obtain deep located areas of myometrium,
- resectobiopsy is characterized by unavailability to obtain deep located areas of myometrium,

high side thermal necrosis, making 70% of preparation impossible for histological analysis.

- Sensitivity of transcervical puncture biopsy for 5-nodular is 48%, 6-8 nodular – 83%, this method provides possibility to obtain deep located areas of myometrium;
- transabdominal puncture biopsy gets tissue samples from external zones of myometrium, making impossible determine the depth of endometrial invasion. Sensitivity of transabdominal biopsy is 58% for 8-nodular biopsy.

Conclusions. 6-nodular transcervical biopsy – is the optimal method to obtain histological samples confirming adenomyosis.

Povzun S.A.¹, Fridman D.B.^{1,2}

¹ Department of Pathomorphology,

² Department of Obstetrics and Gynecology Military- medical academy, Saint Petersburg, Russia

THE CORRELATION OF HISTOLOGICAL AND ULTRASONIC SIGNS OF ADENOMYOSIS

Introduction. Ultrasonography (USI) is a routine investigation technique in gynecology. USI – is the only method, which allows to suspect adenomyosis without invasive study. The determining of US features correlation with endometriometrial alteration helps to define specificity and reliability of different US signs.

Material and methods. USI and histological examination findings of 24 patients with urgent nongravid metrorrhage were observed.

Results. Small cystical inclusions in myometrium can't be explained by heterotopy presence, but connected with

fibroid changes of myometrium, and besides the effectiveness of sign reduces while the vascular component rises, because of the diminution of specific volume of fibroid, and inability to recognize separate inclusions in case of diffuse abnormal echostructure of myometrium. Diffuse echostructure discovered at USI is related to muscular tissue edema, most evident at paraheterotopic areas of myometrium. Most effective USI sign in adenomyosis diagnosis was – M-echo visualization at third day after uterine curettage, which reflects basal layer of endometrium and adjacent myometrium hyperplasia.

Popov A.A., Machanskite O.V., Manannikova T.N.,
Shaginyn G.G., Ramazanov M.R.

Dp. Endoscopy Moscow Reg. Research Institute, Moscow, Russia

THE RATE OF PREGNANCY FOLLOWING DIFFERENT TREATMENT OF ENDOMETRIOSIS

Materials and methods. Endometriosis surgery was carried out in 401 patients. Average age was 27,8 and all of them suffered from infertility for 2-5 years while endocrine factor was excluded. Patients were divided into two groups: a) surgical treatment only b) surgical treatment followed by hormonal therapy Gn-RH agonists for 6 months.

Results. Pregnancy rate in patients with endometri-

osis associated infertility reached 45,2%. It was noted that pregnancy frequency in patients undergoing combined treatment was 18% higher that in patients which underwent only surgical destruction of the nidi. Majority of pregnancies occurred in the first six months following the treatment, after that pregnancy frequency sharply declined.

Conclusion. Two stage treatment (1 – destruction of

the endometriosis nodi, 2 – pharmacotherapy) is pathomorphologically substantiated. First stage envisages destruction of all or most of endometriosis nodi which cause num-

ber of condition leading to depressed generative function, while the second stage inactivates eutopic endometrium which is the constant source of endometriosis.

Ruhliada N.N.

SPb scientific research institute of emergency medicine by I.I.Dzhanelidze, St.-Petersburg, Russia

Preface. Hormonal therapy is the integral component of complex treatment of all forms of endometriosis. The saved up 20-years world experience shows, that the best results are achieved with GnRH-agonists using, and the basic mechanism of their action is amenorrhea achievement. The mentioned mono- or adjuvant therapy is always coexisting with development of by-effects: the developed picture of climacteric infringements, mental and vegetative frustration. It does impossible and inexpedient carrying out of hormonal therapy during more, than 6-8 months. Frequency of relapse of clinical signs reaches 48% after GnRH canceling within first two years of supervision. Works by Thipgen J.T. et al. (2004) have proved expediency of carrying out of intraperitoneal chemotherapy with doxorubicin and cisplatin in disseminated endometrial cancer management, and also have confirmed efficiency of application intraperitoneal chemotherapy (ICT) with these drugs with the purpose of suppression of growth and resorption activation of endometrial deposits.

Objective. To show expediency of carrying out of ICT with cytostatics in treatment of the widespread endometriosis.

Ruhliada N.N., Belyaev A.M.

SPb scientific research institute of emergency medicine by I.I.Dzhanelidze, St.-Petersburg, Russia

At a hyperthermal variant it is better to begin the intraperitoneal chemotherapy (ICT) right after performance of a surgical stage. Perfusion is better to carry out by introduction through troacar ports of 2 laparoscopic irrigation tips connected to perfusion system. It's need to provide a horizontal position of the patient, so that the pelvis would be covered with perfusing fluid. An irrigation tip leaves in Douglas pouch, aspirational – collects perfusate from utero-vesical pouch. Perfusion speed is not less than 500 ml/min, perfusing temperature on input – 42°C, duration of perfusion – 30 minutes. The total perfusate amount is 2000-3000 ml in which the

SUBSTANTIATION OF ADJUVANT INTRAPERITONEAL CHEMOTHERAPY IN COMPLEX TREATMENT OF WIDESPREAD ENDOMETRIOSIS (III-IV STAGES)

Material and methods. Carrying out of hyperthermal intraoperative intraperitoneal (ICT) and normothermal postoperative chemotherapy with doxorubicin 60 mg/m² and cisplatin 50 mg/m² at 23 patients with endometriosis III-IV (ASRM staging). Control group – 10 women – adjuvant therapy with GnRH agonists for 3 months

Results. Adjuvant ICT was effective at treatment of the widespread endometriosis and has led to proof positive clinical effect at 93,4% of women within the first year of supervision. Single-staged adjuvant ICT at the widespread endometriosis is not accompanied by development after operation of any by-effects; and performance of second-look laparoscopy at 14 (60,9%) women has shown, that efficiency of ICT is higher, than adjuvant 3-4 month courses of GnRH agonists (Zoladex, Buserelin) and leads to full resorption of endometriosis heterotopies and surrounding infiltration. The use of "soft" modes of CT (30 minutes perfusion of 42°C solutions) was not accompanied by development of adhesive process after operation in any case, and, hence, had no negative mechanical action on fertility patients. The specified variant of therapy did not result in ovulatory function disturbances.

OPTIMAL PROTOCOL OF NORMO- AND HYPERTHERMAL INTRAPERITONEAL CHEMOTHERAPY (ICT) IN COMPLEX MANAGEMENT OF THE WIDESPREAD ENDOMETRIOSIS (III-IV STAGES)

demanding dose of cytostatics is contained. There could be a fractional injection of 1000 ml of warm (42°C) perfusate intraabdominally and its replacement each 3-4 minutes with new portion of warm solution – duration has to be increased up to 40 minutes, we leave drainage for removal of the rests of perfusate.

At normothermal variant – after performance of a surgical stage we place in an abdominal cavity 1000 ml of the saline containing cytostatics dose. In Douglas pouch we leave a drainage not less than 7 mm in diameter which was opened later with 20-24 hours after the primary surgery for removal of the rests of perfusing fluid. Calculation of cy-