

ВАРИАНТЫ ВРЕМЕННОЙ ИММОБИЛИЗАЦИИ ПРИ ПЕРЕЛОМАХ ЧЕЛЮСТЕЙ

© Н.Е. Митин, Т.С. Родина, Н.Н. Стрелков, М.И. Золотова, В.В. Волкова

ФГБОУ ВО Рязанский государственный медицинский университет
им. акад. И.П. Павлова Минздрава России, Рязань, Россия

Как в мирное, так и в военное время переломы костей лицевого скелета являются актуальным явлением (от 5-6% до 7-9% от всех травматических повреждений скелета) и требуют рациональных методов лечения. При этом, на долю переломов нижней челюсти приходится 65-85%, верхней челюсти – 4-6%, скуловой и носовой костей – 7-9% и 4-7% соответственно. Необходимость иммобилизации обусловлена риском возникновения болевых ощущений и осложнений, таких как кровотечение, микробная контаминация, асфиксия, повреждение нервов и сосудов. В статье проводится сравнительный анализ способов транспортной иммобилизации челюстей на примере травм, полученных в условиях военного времени или в момент чрезвычайных ситуаций. Были выбраны внеротовые методы, удерживающие отломки челюстей посредством повязки, прикрепленной к мозговой части черепа, и внутриротовые, имеющие межчелюстное лигатурное скрепление. **Заключение.** Стандартная пращевидная повязка Энтина имеет преимущество по сравнению с повязками по Гиппократу и по Померанцевой-Урбанской. Она достаточно универсальна и может быть подогнана под размер головы любого пострадавшего, обладает приемлемой прочностью и надежностью фиксации челюстей. Выявлены преимущества методики Айви в сравнении с другими методами межчелюстного лигатурного скрепления. Ее достоинством является прочная и качественная иммобилизация отломков челюстей, возможность использования при множественных переломах, простота применения, что позволяет рекомендовать методику Айви для использования при возникновении показаний.

Ключевые слова: перелом челюсти, временная иммобилизация, повязка, лигатурное скрепление.

VARIANTS OF TEMPORARY IMMOBILIZATION IN JAW FRACTURES

N.E. Mitin, T.S. Rodina, N.N. Strelkov, M.I. Zolotova, V.V. Volkova

Ryazan State Medical University, Ryazan, Russia

In peacetime and in wartime, fractures of the bones of the facial skeleton are an important phenomenon (5-6% to 7-9% of all traumatic skeletal injuries) requiring use of rational treatment methods. The share of mandibular fractures accounts for 65-85%, of maxilla – 4-6%, of zygomatic and nasal bones – 7-9% and 4-7%, respectively. The need for immobilization is associated with the risk of development of pain and complications, such as bleeding, microbial contamination, asphyxia, injury of nerves and blood vessels. The article gives a comparative analysis of means of transport immobilization of jaws on examples of traumas received in the wartime and in emergen-



cy situations. There were chosen extraoral methods that hold jaw fragments with the help of bandage secured to the cerebral cranium, and intraoral methods with maxillomandibular ligature fixation. **Conclusion.** Standard Entin head-chin strap has certain advantages over Hippocratic cap bandage and Pomerantseva-Urbanskaya method. It is sufficiently universal and may be adjusted to any size of the head, possesses sufficient strength and provides reliable fixation of jaws. Ivy's method was found to have advantages in comparison with other methods of maxillomandibular ligature fixation, which are strong and high-quality immobilization of the jaw fragments, possibility of use with multiple fractures, simplicity of application. All this permits to recommend Ivy's method for use in case of indications.

Keywords: *jaw fracture, temporary immobilization, bandage, ligature fixation.*

In conditions of wartime and in emergency situations, the first medical assistance is given at the stage of evacuation. Temporary immobilization of fractures is of much importance in this period due to a risk of development of complications in the maxillofacial region threatening the life of the victim. There often arises a question of selection of a rational method of transport immobilization of jaws based on the peculiarities of the clinical picture of the trauma.

Thus, the *aim* of work was comparison of methods and means of temporary immobilization of the jaws based on the analysis of literature sources.

For immobilization of jaws transportation splints are used that are bandages of different design. In the literature methods of use of bandages are described for fixation of jaw fractures with characteristic clinical peculiarities [1-3]. To arrest bleeding in case of fracture of the upper jaw and lower jaw beyond the dental arch, compressive bandage is often applied. In case of jaw fractures under field conditions with multiple ruptures of soft tissues, a variant of standard Entin's head-chin strap is considered for use [4]. The choice of this method was determined by existence of this fixture in a standard tool bag of a medical instructor of Russian Armed Forces and Ministry of Emergency Situation of Russia. According to T.P. Dyubkova, in transportation of the victim with use of Entin's strap no purulo-inflammatory complications occur on the 3rd-

4th day, and the bone fragments of jaws are reliably fixed. The author notes that in fractures of the lower jaw within the dental arch, this bandage should be used only to hold the bone fractures, since excessive pressure on the fragments may worsen their displacement and cause asphyxia [5].

Use of Hippocratic cap bandage in combined fracture of the upper and lower jaw was considered an alternative to Entin's strap, since it may be applied by a specialist of any qualification. In case of fracture of both jaws it holds the bone fragments and prevents their displacement.

Variants of the extraoral temporary immobilization were considered on an example of Hippocratic cap in comparison with other methods. L.M. Marmysheva, et al. experimentally determined dependence of the time of development of purulo-inflammatory complications on speed of application of the bandage. The authors think that the faster the bandage is applied, the lower the probability for development of complications in the maxillofacial region [6]. I.S. Kopetsky states that use of these structures does not require any invasive manipulations for their application [7]. According to many authors, risk of aspiration in case of vomiting may be eliminated by a rapid removal of the bandage by a victim without any effort. However, Hippocratic head-chin bandage does not satisfy all the requirements: immobilization of the jaw fragments is insufficiently rigid, and this ban-

dage made of bandage gauze, is unreliably secured to the head and often slips down especially in loss of consciousness [8-10].

Some authors give preference to a standard elastic four-tailed bandage of Pomerantseva-Urbanskaya [11-13]. In experiments, this bandage proved to be simple in use and convenient, that is, its application does not require much effort. V.Yu. Ziambetov notes impossibility of using this bandage in fractures of edentulous jaws [14].

The majority of authors in their works consider use of means that may be at hand (a pen, a stick, a plate) for temporary immobilization secured to the head with a handkerchief, rope, belt, and serving an improvised fixating base [15,16]. K.A. Sveshnikov proposes use of the undamaged removable dentures together with head-chin strap as a means of transportation immobilization in case of fractures of edentulous jaws [17].

Fixation of fragments of the low jaw within the dental arch, fractures of the alveolar process of the upper jaw and of multiple fractures of jaws can be achieved using maxillomandibular ligature tying. Some authors use a wire 0.5 mm thick as a ligature [18]. V.V. Maslyakov, et al. found out a common condition for use of all intraoral immobilization methods. They state that for one-jaw bandage the presence of 2-3 stable teeth is required on each fragment of the lower jaw, and for two-jaw bandage there should be stable teeth in both jaws with the undamaged upper jaw. Use of this method is limited in fractures of both jaws and in the presence of mobile teeth or teeth in the fracture zone [19]. I.Yu. Lebedenko gives the priority to Ivy's method of ligature fixation based on the introduction of wire in the form of a hairpin through the interdental space with twisting of its ends in the vestibule of the mouth. The author notes that in case of fracture in the region of a branch or in the zone of the second and third molars of the lower jaw this method is unrealizable, since there is no possibility to

pass the ligature on both sides of the fracture line. In opening the mouth, for example, in vomiting, the vertical wires are readily cut, and the main part of the structure is retained in the mouth cavity [20]. A.V. Lepilin noted that parodontium of teeth that participates in the maxillomandibular fastening, is under the load of holding the jaw fragments in correct position and of counteracting muscles that lower the mandible [21].

For comparison, I.M. Samokhvalov considers the possibility of using Kazanyan's method based on the passing of the ligature around the teeth of jaw fragments and antagonistic teeth in the form of "eight" with twisting of the wire ends in the vestibule of the mouth. The method suggests existence of a thick wire bundle in the vestibule of the mouth which may traumatize oral mucosa. Besides, I.M. Samokhvalov indicates the necessity of repeated application of ligatures in case of their breakage or after untwisting for examination or for hygienic procedures in the oral cavity [22].

The same author emphasizes peculiarities of Vilga's method with use of a wire requiring fixation with special structures in the form of a button, and of Geikin's method using lead pellets for fixation of the ends of the wire in the mouth cavity, which is ecologically unfriendly and difficult for access [23].

A.A. Mikhailov emphasizes method of Gotsko that uses a polyamide thread instead of ligature wire which is tied on the vestibular side and passes through the interdental space from the vestibule into the mouth cavity and back. The author notes that the method is less traumatic and rather effective, but is difficult for implementation in conditions of limited time and without special qualification [24].

M.V. Peshkov, et al. came to the conclusion about unreasonableness of using mobile teeth or teeth in the zone of fracture that are subject to obligatory extraction. If possible, pairs of stable antagonistic teeth should be used [25].

After analysis and comparison of peculiarities of methods of temporary immobilization we think it would be reasonable to present the generalized data in the form of a table (Tab. 1).

Thus, it was ascertained that use of extraoral methods of jaw immobilization has a number of advantages. A possibility of use of bandages in the early stage of medical evacuation prevents development of complications of the maxillofacial region and permits to provide fast and timely fixation of the jaw fragments. The peculiarity of bandages consists in holding of the soft tissue flaps in case of multiple ruptures, and of fragments of edentulous jaws, which is impossible with use of ligatures. Convenience of assembly and disassembly of all extraoral immobilization methods, and also of methods of maxillomandibular ligature fixation by Ivy and Gotsko, is a positive moment under risk of aspiration with vomitus. However, use of bandages is ineffective in multiple fractures of jaws and does not provide sufficient quality of fixation of the fractured jaw fragments (which is the advantage of ligature).

The most commonly used and effective extraoral structure of temporary immobilization of jaws is Entin's standard four-tail bandage that possesses common for all bandages advantages and disadvantages, and, besides, has individual advantages. Existence of this fixture in a standard kit of combat medics of Russian Armed Forces and Ministry of Emergency Situations of Russia, determines its

predominant use in military environment and emergency conditions. Use of rubber cords attached to a standard cap ensures a reliable fixation of jaw fragments, which makes it superior to bandages of other structures: of Hippocrates and Pomerantseva-Urbanskaya.

It was also found that Ivy's method of maxillomandibular ligature fixation possesses the most of advantages: simplicity of structure and convenience of use, high-quality and reliable fixation of jaw fragments, absence of additional fixation elements characteristic of Vilga's and Geikin's methods. Method of Gotsko is difficult in implementation which is a considerable drawback. All intraoral methods are characterized by bad accessibility in the field which considerably limits their application.

Conclusion

On the basis of the obtained results it may be concluded that standard Entin's four-tail bandage possesses advantages over Hippocratic cap and Pomerantseva-Urbanskaya bandage. It is sufficiently universal and may be adjusted to any size of head, possesses adequate strength and provides reliable fixation of jaws.

Ivy's method possesses advantages in comparison with other considered methods of maxillomandibular ligature fixation. Its advantages consist in a strong and reliable immobilization of jaw fragments, possibility of use in case of multiple fractures, simplicity of implementation, which permits to recommend it for application with indications.

Литература

1. Ургуналиев Б.К., Кулназаров А.С. Современное состояние проблем травматологии челюстно-лицевой области (обзор литературы) // Вестник КГМА им. И.К. Ахунбаева. 2016. №3. С. 60-63.
2. Ластовецкий А.Г., Лебедев М.В., Аверьянова Д.А. Частота и структура травматических повреждений мозгового и лицевого отделов черепа у пострадавших в дорожно-транспортных происшествиях // Известия высших учебных заведений. Поволжский регион. Медицинские науки. 2014. Т. 31, №3. С. 105-116.
3. Гуйтер О.С., Митин Н.Е., Устюгова А.Е., и др. Этиологические факторы, способствующие возникновению дефектов и деформаций челюстно-лицевой области // Наука молодых (Eruditio Juvenium). 2015. №4. С. 91-97.
4. Мирошниченко Ю.В., Бояринцев В.В., Бунин С.А., и др. Использование комплектов медицинского имущества, наборов и укладок медицинских для ликвидации медико-санитарных последствий чрезвычайных ситуаций // Медико-биологические и социально-психологические проблемы безопасности в чрезвычайных ситуациях. 2014. №3. С. 39-48.

Table 1

**Comparison of Extraoral and Intraoral Methods
of Temporary Fixation of Jaws**

Method	Assessment Criterion														
		Availability of material in the field	Convenience of assembly and disassembly in mouth cavity	Existence of elastic maxillomandibular ligatures	Existence of rigid maxillomandibular ligatures	Existence of additional fixation elements	Existence of standard fixation cap	Fixation to brain part of skull	Speed of application	Quality of fixation of jaws	Use on edentulous jaws	Use in multiple fractures	Use in multiple ruptures of soft tissues	Use at early stage of medical evacuation	
Extraoral methods (bandages)	Entin	-	+	+	-	+	+								
	Hippocrates	+	+	-	-	-	-	+	+	-	+	-	+	+	
	Pomerantseva-Urbanskya	-	+	+	-	+	+								
Intraoral methods (ligature fixation)	Ivy	-	+	-	+	-	-								
	Kazanyan	-	-	-	+	-	-								
	Vilga	-	-	-	-	+	-	-	-	+	-	+	-	-	
	Geikin	-	-	-	-	+	-								
	Gotsko	-	+	+	-	-	-								

- Дюбкова Т.П. Обоснование необходимости разработки стандартов первой помощи пострадавшим в чрезвычайных ситуациях // Проблемы стандартизации в здравоохранении. 2012. №1-2. С. 29-32.
- Мармышева Л.Н., Осипов А.Б., Цыбенко Е.И., и др. Система аварийно-спасательных и других неотложных работ и их характеристика. В кн.: Безопасность жизнедеятельности. СПб.: Издательство СПбГЭУ; 2016. С. 180-223.
- Копецкий И.С., Насибуллин А.М., Кабисова Г.С. Оказание медицинской помощи больным с огнестрельными ранениями челюстно-лицевой области в условиях многопрофильной больницы // Наука Красноярья. 2012. Т. 1, №3. С. 78-88.
- Копецкий И.С., Насибуллин А.М., Кабисова Г.С., и др. Ранения челюстно-лицевой области огнестрельного характера в условиях мегаполиса // Медицинский вестник МВД. 2011. Т. 55, №6. С. 19-23.

9. Флейшер Г.М. Особенности клинической картины переломов нижней челюсти // Символ науки. 2016. Т. 14, №2-3. С. 178-182.
10. Тимошенко Л.И., Юров И.Б. Первая помощь пострадавшим в дорожно-транспортном происшествии. Ставрополь: АГРУС; 2013.
11. Байриков И.М., Фишер И.И., Шухорова Ю.А., и др. Совершенствование оказания медицинской помощи больным с переломами нижней челюсти на всех этапах реабилитации // Управление качеством медицинской помощи. 2017. №1-2. С. 59-65.
12. Башинский О.А., Гудзь Ю.В., Иванов А.И., и др. Рекомендации по основам оказания первой помощи пострадавшим в чрезвычайных ситуациях сотрудниками, военнослужащими и работниками государственной противопожарной службы и спасателями аварийно-спасательных формирований и аварийно-спасательных служб МЧС России. СПб.: Политехника-сервис; 2015.
13. Поспелов Н.Н., Лукьяненко А.В., Баженов О.В., и др. Особенности организации медицинской помощи раненым в лицо в современных локальных военных конфликтах // Медицинский вестник МВД. 2017. Т. 91, №6. С. 27-31.
14. Зиамбетов В.Ю., Востриков В.А., Зиамбетов Вл.Ю. Основы самообороны и рукопашного боя в профессионально-прикладной физической культуре студентов. Оренбург: Издательство ОГПУ; 2016.
15. Копецкий И.С., Насибуллин А.М., Гончарова А.В. Огнестрельные ранения челюстно-лицевой области мирного времени // Вестник Российского государственного медицинского университета. 2012. №1. С. 25-29.
16. Мишин Ю.А., Забродский Д.С., Меркулов А.В. Совершенствование комплектно-табельного оснащения для оказания первой помощи в военноморском флоте // Морская медицина. 2015. Т. 1, №2. С. 55-57.
17. Свешников К.А., Якушев Д.Б. Транспортная имобилизация на этапе скорой помощи // Справочник фельдшера и акушерки. 2014. №10. С. 26-32.
18. Севбитов А.В., Митин Н.Е., Браго А.С., и др. Основы зубопротезной техники. Ростов н/Д: Феникс; 2016.
19. Масляков В.В., Барсуков В.Г., Куркин К.Г., и др. Объем оказания первичной доврачебной медико-санитарной помощи лицам из населения с огнестрельными ранениями в условиях локального вооруженного конфликта // Медицина катастроф. 2018. Т. 102, №2. С. 30-33.
20. Лебедево И.Ю., Каливрадзян Э.С. Ортопедическая стоматология [Электронный ресурс]. М.: ГЭОТАР-Медиа; 2011.
21. Лепилин А.В., Ерокина Н.Л., Захарова Н.Б., и др. Влияние различных методов иммобилизации на ткани пародонта у больных хроническим генерализованным пародонтитом при переломах нижней челюсти // Российский стоматологический журнал. 2011. №3. С. 25-28.
22. Самохвалов И.М., Рева В.А. Догоспитальная помощь раненым в военных конфликтах: состояние и перспективы // Военно-медицинский журнал. 2015. Т. 336, №10. С. 15-26.
23. Самохвалов И.М., Казначеев М.В., Петухов К.В., и др. Догоспитальная сортировка как первый этап управления лечением пострадавших // Скорая медицинская помощь. 2016. Т. 17, №2. С. 13-19.
24. Михайлов А.А., Ботыгин В.И., Гинко В.И., и др. Основы начальной военной подготовки. Шуя: Издательство Шуйского филиала Ивановского государственного университета; 2015.
25. Пешков М.В., Гуревич К.Г. Социальная характеристика и качество жизни пациентов, обращающихся за стоматологической хирургической помощью // Российский медико-биологический вестник имени академика И.П. Павлова. 2016. Т. 24, №1. С. 123-128.

References

1. Urgunaliyev BK, Kulnazarov AS. Traumatology of the maxillofacial region: current condition of the problems (Literature review). *Vestnik KSMA named after I.K. Akhunbaev*. 2016;3:60-3. (In Russ).
2. Lastoveckiy AG, Lebedev MV, Aver'yanova DA. Frequency and structure of traumatic damages of brain and facial part of skull in road traffic accident victims. *University Proceedings. Volga Region. Medical Sciences*. 2014;31(3):105-16. (In Russ).
3. Guyter OS, Mitin NE, Ustyugova AE, et al. Etiological factors promoting appearance of defects and deformations of maxillofacial area. *Science of the Young (Eruditio Juvenium)*. 2015;4:91-7. (In Russ).
4. Miroshnichenko YV, Boyarincev VV, Bunin SA, et al. Use of medical sets and kits during elimination of health consequences of emergencies. *Medico-Biological and Socio-Psychological Problems of Safety in Emergency Situations*. 2014;3:39-48. (In Russ).
5. Dyubkova TP. The substantiation for necessity of working out of the primary help standards for victims of emergency situation. *Problemy Standartizacii v Zdravoohranenii*. 2012;1-2:29-32. (In Russ).
6. Marmysheva LN, Osipov AB, Tsybenko EI, et al. Sistema avariyno-spatel'nyh i drugih neotlozhnyh rabot i ih harakteristika. In: *Bezopasnost' zhiznedeyatel'nosti*. Saint Petersburg: Izdatel'stvo SPbGEU; 2016:180-223. (In Russ).
7. Kopeckiy IS, Nasibullin AM, Kabisova GS. Rendering medical care to patients with gunshot facial wounds in the hospital. *Siberian Journal of Economics and Management*. 2012;1(3):78-88. (In Russ).
8. Kopeckiy IS, Nasibullin AM, Kabisova GS, et al. Maxillofacial gunshot wounds in the context of a

- megalopolis. *MIA Medical Bulletin*. 2011;55(6):19-23. (In Russ).
9. Fleysher GM. Osobennosti klinicheskoy kartiny perelomov nizhney chelyusti. *Simvol Nauki*. 2016; 14(2-3):178-82. (In Russ).
 10. Timoshenko LI, Yurov IB. *Pervaya Pomoshch' Postradavshim v Dorozhno-transportnom Proisshествii*. Stavropol': AGRUS; 2013. (In Russ).
 11. Bayrikov IM, Fisher II, Shukhorova YuA, et al. Improvement of medical care to patients with jaw fractures at all stages of rehabilitation. *Upravlenie Kachestvom Medicinskoj Pomoshchi*. 2017;1-2:59-65. (In Russ).
 12. Bashinskiy OA, Gud' YuV, Ivanov AI, et al. *Rekomendacii po osnovam okazaniya pervoy pomoshchi postradavshim v chrezvychaynykh situatsiyah sotrudnikami, voennosluzhashchimi i rabotnikami gosudarstvennoy protivopozharnoy sluzhby i spasatelyami avariyno-spasatel'nykh formirovaniy i avariyno-spasatel'nykh sluzhb MCHS Rossii*. Saint Petersburg: Politehnika-servis; 2015. (In Russ).
 13. Pospelov NN, Luk'yanenko AV, Bazhenov OV, et al. Specifics of organization of medical care for casualties with maxillofacial wounds in present local military conflicts. *MIA Medical Bulletin*. 2017; 91(6):27-31. (In Russ).
 14. Ziambetov VYu, Vostrikov VA, Ziambetov VIYu. *Osnovy samooborony i rukopashnogo boya v professional'no-prikladnoy fizicheskoy kul'ture studentov*. Orenburg: Izdatel'stvo OGPU; 2016. (In Russ).
 15. Kopetskiy IS, Nasibullin AM, Goncharova AV. Gunshot wounds of the midface in peacetime. *Bulletin of Russian State Medical University*. 2012;1:25-9. (In Russ).
 16. Mishin YuA, Zabrodskiy DS, Merkulov AV. Improvement of complete and organic equipment for first-aid treatment in navy. *Marine Medicine*. 2015; 1(2):55-7. (In Russ).
 17. Sveshnikov KA, Yakushev DB. Transportnaya immobilizatsiya na ehtape skoroy pomoshchi. *Spravochnik Fel'dshera i Akusherki*. 2014;10:26-32. (In Russ).
 18. Sevbitov AV, Mitin NE, Brago AS, et al. *Osnovy zuboproteznoy tekhniki*. Rostov-on-Don: Feniks; 2016. (In Russ).
 19. Maslyakov VV, Barsukov VG, Kurkin KG, et al. Scope of Provision of Primary Medical and Sanitary Care to Members of Population with Gunshot Wounds in Local Armed Conflict Environment. *Disaster Medicine*. 2018;102(2):30-3. (In Russ).
 20. Lebedenko IYu, Kalivradzhiyan EhS. *Ortopedicheskaya stomatologiya* [Electronic resource]. Moscow: GEOTAR-Media; 2011.
 21. Lepilin AV, Erokina NL, Zakharova NB, et al. The influence of various immobilization techniques on periodontal tissue in the patients with mandibular fractures presenting with chronic generalized periodontitis. *Russian Journal of Dentistry*. 2011;3:25-8. (In Russ).
 22. Samokhvalov IM, Reva VA. Dogospital'naya pomoshch' ranenym v voennykh konfliktakh: sostoyanie i perspektivy. *Voенно-Medicinskiy Zhurnal*. 2015; 336(10):15-26. (In Russ).
 23. Samokhvalov IM, Kaznacheyev MV, Petukhov KV, et al. Prehospital triage as the first period of injured patients care management. *Emergency Medical Care*. 2016;17(2):13-9. (In Russ).
 24. Mikhaylov AA, Botygin VI, Ginko VI, et al. *Osnovy nachal'noy voennoy podgotovki*. Shuya: Izdatel'stvo Shuyskogo filiala Ivanovskogo gosudarstvennogo universiteta; 2015. (In Russ).
 25. Peshkov MV, Gurevich KG. Social characteristics and quality of life of patients seeking surgical care dental. *I.P. Pavlov Russian Medical Biological Herald*. 2016;24(1):123-8. (In Russ).

Дополнительная информация [Additional Info]

Источник финансирования. Бюджет ФГБОУ ВО Рязанский государственный медицинский университет им. акад. И.П. Павлова Минздрава России. [Financing of study. Budget of Ryazan State Medical University.]

Конфликт интересов. Авторы декларируют отсутствие явных и потенциальных конфликтов интересов, о которых необходимо сообщить, в связи с публикацией данной статьи. [Conflict of interests. The authors declare no actual and potential conflict of interests which should be stated in connection with publication of the article.]

Участие авторов. Митин Н.Е. – концепция работы, редактирование, Родина Т.С., Стрелков Н.Н., Волкова В.В. – написание текста, редактирование, Золотова М.И. – сбор материала, перевод. [Participation of authors. N.E. Mitin – concept of work, editing, T.S. Rodina, N.N. Strelkov, V.V. Volkova – writing the text, editing, M.I. Zolotova – collections of material, translation.]

Информация об авторах [Authors Info]

Митин Николай Евгеньевич – к.м.н., доцент, заведующий кафедры ортопедической стоматологии и ортодонтии с курсом пропедевтики стоматологических заболеваний ФГБОУ ВО Рязанский государственный медицинский университет им. акад. И.П. Павлова Минздрава России, Рязань, Россия. [**Nikolay E. Mitin** – MD, PhD, Associate Professor, Head of the Department of Orthopedic Dentistry and Orthodontics with the Course of Propaedeutics of Dental Diseases, Ryazan State Medical University, Ryazan, Russia.]

ORCID ID: 0000-0001-7890-6353.

Родина Татьяна Сергеевна – к.м.н., доцент кафедры ортопедической стоматологии и ортодонтии с курсом пропедевтики стоматологических заболеваний ФГБОУ ВО Рязанский государственный медицинский университет им. акад. И.П. Павлова Минздрава России, Рязань, Россия. [**Tat'yana S. Rodina** – MD, PhD, Associate Professor of the Department of Orthopedic Dentistry and Orthodontics with the Course of Propaedeutics of Dental Diseases, Ryazan State Medical University, Ryazan, Russia.]

SPIN: 5723-3633, ORCID ID: 0000-0002-7678-0029, Researcher ID: V-1963-2018.

Стрелков Николай Николаевич – к.м.н., ассистент кафедры ортопедической стоматологии и ортодонтии с курсом пропедевтики стоматологических заболеваний ФГБОУ ВО Рязанский государственный медицинский университет им. акад. И.П. Павлова Минздрава России, Рязань, Россия. [**Nikolay N. Strelkov** – MD, PhD, Assistant of the Department of Orthopedic Dentistry and Orthodontics with the Course of Propaedeutics of Dental Diseases, Ryazan State Medical University, Ryazan, Russia.]

SPIN: 6421-8603, ORCID ID: 0000-0001-9376-3952, Researcher ID: U-9716-2018.

Золотова Мария Игоревна – аспирант кафедры ортопедической стоматологии и ортодонтии с курсом пропедевтики стоматологических заболеваний ФГБОУ ВО Рязанский государственный медицинский университет им. акад. И.П. Павлова Минздрава России, Рязань, Россия. [**Mariya I. Zolotova** – PhD Student of the Department of Orthopedic Dentistry and Orthodontics with the Course of Propaedeutics of Dental Diseases, Ryazan State Medical University, Ryazan, Russia.]

SPIN: 9320-1167, ORCID ID: 0000-0003-3933-9012, Researcher ID: V-2628-2018.

* **Волкова Василиса Вадимовна** – клинический ординатор кафедры ортопедической стоматологии и ортодонтии с курсом пропедевтики стоматологических заболеваний ФГБОУ ВО Рязанский государственный медицинский университет им. акад. И.П. Павлова Минздрава России, Рязань, Россия. [**Vasilisa V. Volkova** – Clinical Resident of the Department of Orthopedic Dentistry and Orthodontics with the Course of Propaedeutics of Dental Diseases, Ryazan State Medical University, Ryazan, Russia.]

SPIN: 8531-6106, ORCID ID: 0000-0002-7619-7597, Researcher ID: U-7506-2018. E-mail: Vasiliska101@mail.ru

Цитировать: Митин Н.Е., Родина Т.С., Стрелков Н.Н., Золотова М.И., Волкова В.В. Варианты временной иммобилизации при переломах челюстей // Российский медико-биологический вестник имени академика И.П. Павлова. 2018. Т. 26, №4. С. 559-566. doi:10.23888/PAVLOVJ2018264559-566

To cite this article: Mitin NE, Rodina TS, Strelkov NN, Zolotova MI, Volkova VV. Variants of temporary immobilization in jaw fractures. *I.P. Pavlov Russian Medical Biological Herald*. 2018;26(4):559-66. doi:10.23888/PAVLOVJ2018264559-566

Поступила/Received: 21.09.2018

Принята в печать/Accepted: 12.12.2018